

BONUS DIGITAL CONTENT

Adult Preventive Health Care Schedule: Recommendations from the USPSTF (as of May 9, 2023)

To be used in conjunction with USPSTF recommendation statements (for additional details see tables and references at <https://www.aafp.org/afp/PHCS>)

Only grade A/B recommendations are shown

Age	18	21	24	25	35	40	45	50	55	59	65	70	74	75	80
USPSTF screening recommendations															
Alcohol misuse ¹	(B)														
Unhealthy drug use ²	(B)														
Depression ³	(B)														
Hypertension ⁴	(A)														
Obesity/weight loss ⁵	(B) if BMI 30 kg per m ² or greater														
Tobacco use and cessation ⁶	(A)														
Hepatitis C virus infection ⁷	(B)														
HIV infection ⁸	(A) (A) if at increased risk														
Hepatitis B virus infection ⁹	(B) if at increased risk														
Syphilis ¹⁰	(A) if at increased risk														
Latent tuberculosis infection ¹¹	(B) if at increased risk														
BRCA gene risk assessment ¹²	(B) if appropriate personal or family history of BRCA-related cancer or ancestry														
Chlamydia and gonorrhea ¹³	(B) if sexually active (B) if at increased risk														
Intimate partner violence ¹⁴	(B) women of childbearing age														
Cervical cancer ¹⁵	(A) See p. 3 for test options and screening intervals														
Prediabetes and type 2 diabetes ¹⁶	(B) if overweight or obese														
Colorectal cancer ¹⁷	(B) (A)														
Breast cancer ¹⁸	(B) biennial screening														
Lung cancer ¹⁹	(B) if 20-pack-year history and current or former smoker (quit in past 15 years)														
Osteoporosis ²⁰	(B) if postmenopausal and elevated risk (B)														
Abdominal aortic aneurysm ²¹	(B) if an "ever smoker"														

USPSTF preventive therapies recommendations

HIV preexposure prophylaxis ²²	(A) if at high risk of HIV infection														
Primary prevention of breast cancer ²³	(B) offer if at increased risk for breast cancer and low risk for side effects														
Folic acid supplementation ²⁴	(A) if capable of conceiving														
Statins for primary prevention of CVD ²⁵	(B) see criteria on p. 4														
Fall prevention in community-dwelling older adults ²⁶	(B) exercise interventions if at increased fall risk														

USPSTF counseling recommendations

Sexually transmitted infection prevention ²⁷	(B) if at increased risk														
Diet/activity for CVD prevention ²⁸	(B) adults with CVD risk factors														
Skin cancer prevention ²⁹	(B) if fair skinned														
Healthy weight gain in pregnancy ³⁰	(B) all pregnant women														

Legend

	Normal risk	With specific risk factor	Recommendation grades
Recommendation for men and women			A Recommended (likely significant benefit)
Recommendation for men only			B Recommended (likely moderate benefit)
Recommendation for women only			C Do not use routinely (benefit is likely small)
			D Recommended against (likely harm or no benefit)
			I Insufficient evidence to recommend for or against

BMI = body mass index; CVD = cardiovascular disease; USPSTF = U.S. Preventive Services Task Force.

Visual adaptation from recommendation statements by Swenson PF, Lindberg C, Carrilo C, and Clutter J.

HIV RISK FACTORS

- | | |
|----------------------------------|---|
| IV drug use | Sex with individuals who are IV drug users, bisexual, or HIV positive |
| Men who have sex with men | |
| Other STI | Unprotected sex, including anal intercourse |
| Requesting STI testing | |
| Sex exchanged for drugs or money | |

Patients in whom to consider PrEP:

Sexually active men who have sex with men who have any of the following:

- Sexual relationship with serodiscordant partner
- Inconsistent use of condoms during anal sex
- Syphilis, gonorrhea, or chlamydia infection in past six months

Sexually active heterosexual patients with any of the following:

- Sexual relationship with serodiscordant partner
- Inconsistent use of condoms with high-risk partner
- Syphilis or gonorrhea infection in past six months

Injection drug users with any of the following:

- Shared drug-injection equipment
- Risks of infection through sex (see above)

IV = intravenous; PrEP = preexposure prophylaxis; STI = sexually transmitted infection.

HEPATITIS B INFECTION RISK FACTORS

- | | |
|------------------------------------|---|
| HIV infection | Men who have sex with men |
| Infected sex partner | Origin from regions* with prevalence \geq 2% |
| Intravenous drug use | U.S.-born children of immigrants from regions* with prevalence \geq 8%, if unvaccinated |
| Living with an infected individual | |

*—Risk of regions can be found at <http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5708a1.htm>.

SYPHILIS RISK FACTORS

- | | |
|----------------------------|----------------------------------|
| High-risk sexual behaviors | Men who have sex with men |
| Incarceration | Sex exchanged for drugs or money |
| Local prevalence | |

TUBERCULOSIS RISK FACTORS

- | | |
|--------------------------------|--|
| Health professionals* | Prisoners, including former |
| Homelessness, including former | Residents of high-risk regions, including former |
| Immunosuppression* | |

*—Evidence for screening not reviewed by the USPSTF because this is standard practice in public health and standard of care for patients with immunosuppression, respectively.

CHLAMYDIA AND GONORRHEA RISK FACTORS

- | | |
|---|--|
| New or multiple sex partners | Sex exchanged for drugs or money |
| Other STI, including history of STI | |
| Partner with STI | Unprotected sex or inconsistent condom use |
| Partners who have multiple sex partners | |

STI = sexually transmitted infection.

CARDIOVASCULAR DISEASE RISK FACTORS

- | | |
|---|---|
| Atherosclerotic cardiovascular disease risk \geq 7.5% | Hypertension or elevated blood pressure |
| Dyslipidemia | Metabolic syndrome |

BREAST CANCER RISK FACTORS

Consider use of a risk-assessment model for patients with a history of biopsy or positive family history

SEXUALLY TRANSMITTED INFECTION RISK FACTORS

Similar to those risk factors listed previously for sexually transmitted infections; consider local and population-based prevalence in individual risk assessment

Adult Preventive Health Care Schedule: Recommendations from the USPSTF

Grade A/B Recommendations (with Associated Grade C/D/I Recommendations):

Alcohol misuse screening¹

(B) Screen adults and provide brief behavioral interventions for risky alcohol use

Unhealthy Drug Use Screening²

(B) Screen all adults older than 18 years for unhealthy drug use (by asking questions, not biological specimens)

Depression screening³

(B) Screen adults with systems for evaluation and management

Hypertension screening⁴

(A) Screen adults; exclude white coat hypertension before starting therapy

Obesity/weight loss screening⁵

(B) Refer adults with obesity to intensive behavioral interventions for weight loss

Tobacco use and cessation screening⁶

(A) Screen all nonpregnant adults and provide behavior therapy and U.S. Food and Drug Administration–approved intervention therapy for cessation

(A) Screen all pregnant women and provide behavior therapy.

(I) IETRFOA electronic nicotine delivery systems for tobacco cessation

(I) Pharmacotherapy for tobacco cessation in pregnant persons

Hepatitis C virus infection screening⁷

(B) Screen adults 18 to 79 years of age

HIV infection screening⁸

(A) Screen individuals 15 to 65 years of age

(A) Screen older and younger persons who are at increased risk

Hepatitis B virus infection screening⁹

(B) Screen adolescents and adults at high risk

Syphilis screening¹⁰

(A) Screen individuals at increased risk

Latent tuberculosis infection screening¹¹

(B) Screen individuals at increased risk

BRCA-related cancer risk assessment/screening¹²

(B) Use a familial risk assessment tool (evaluated assessment tools listed in full text) in women with either:

- Personal or family history of breast, ovarian, tubal, or peritoneal cancers
- Ashkenazi Jewish ancestry (i.e., ancestry with increased risk of *BRCA* mutation)

For positive risk tools, offer genetic counseling and genetic testing, if indicated.

(D) Recommend against screening for patients without appropriate family history, personal history, or ancestry

Chlamydia and gonorrhea screening¹³

(B) Screen sexually active women 24 years and younger, and women at increased risk who are 25 years and older

(I) IETRFOA screening sexually active men

Intimate partner violence screening¹⁴

(B) Screen women of childbearing age and refer to appropriate services

(I) IETRFOA screening all vulnerable and older adults for abuse or neglect

Cervical cancer screening¹⁵

(A) Screen women

- 21 to 29 years of age every three years with cytology alone
- Frequency of screening may increase to every five years for women 30 to 65 years of age with cytology and high-risk human papillomavirus cotesting or high-risk human papillomavirus testing alone

(D) Recommend against screening in women

- 20 years and younger
- Older than 65 years if adequately screened previously and no increased risk of cervical cancer
- With hysterectomy (including cervix) without history of cervical intraepithelial neoplasia grade 2 or 3 or cervical cancer
- Younger than 30 years with human papillomavirus testing alone or in combination with cytology

Prediabetes and type 2 diabetes screening¹⁶

(B) Screen adults 35 to 70 years of age who are overweight or obese and refer patients with abnormal glucose levels for intensive counseling for healthy diet and exercise

Colorectal cancer screening¹⁷

(A) Screen patients 50 to 75 years of age with fecal occult blood (or immunochemical) test, sigmoidoscopy, colonoscopy, computed tomography colonography, or multitargeted stool DNA test

(B) Screen patients 45 to 49 years of age with fecal occult blood (or immunochemical) test, sigmoidoscopy, colonoscopy, computed tomography colonography, or multitargeted stool DNA test

(C) Selectively offer screening to patients 76 to 85 years of age

Breast cancer screening¹⁸

(B) Biennial screening mammography in women 50 to 74 years of age

(C) Screening is an individualized decision for women 40 to 49 years of age

(I) IETRFOA

- Mammography after 75 years of age
- Screening with digital breast tomosynthesis
- Adjunctive screening in women with dense breast tissue and negative screening mammogram

Lung cancer screening¹⁹

(B) Screen annually with low-dose computed tomography for individuals 50 to 80 years of age with a 20-pack-year history who currently smoke or quit within the past 15 years; discontinue screening once a person has not smoked for 15 years or develops a health problem that limits life expectancy

Osteoporosis screening²⁰

(B) Screen women 65 years and older

(B) Screen postmenopausal women if increased fracture risk shown with an osteoporosis risk tool (e.g., 8.4% in 10 years by U.S. FRAX tool)

(I) IETRFOA screening men

continues

CHD = coronary heart disease; CVD = cardiovascular disease; FRAX = Fracture Risk Assessment; IETRFOA = insufficient evidence to recommend for or against; PrEP = preexposure prophylaxis; USPSTF = U.S. Preventive Services Task Force.

Adult Preventive Health Care Schedule: Recommendations from the USPSTF *(continued)*

Grade A/B Recommendations (with Associated Grade C/D/I Recommendations): *(continued)*

Abdominal aortic aneurysm screening²¹

- (B) Screen men 65 to 75 years of age who ever smoked (100 or greater lifetime cigarettes) with one-time abdominal aortic aneurysm ultrasonography
- (C) Recommend selective screening of men 65 to 75 years who have never smoked
- (I) IETRFOA women 65 to 75 years of age who ever smoked
- (D) Recommend against routine screening in women 65 to 75 years of age who have never smoked

HIV prevention with PrEP²²

- (A) Offer PrEP to persons at high risk of infection. See original text for considerations in patient selection

Primary prevention of breast cancer²³

- (B) Consider medications (such as tamoxifen, raloxifene, or aromatase inhibitors) that reduce risk of breast cancer in women at increased risk though with low risk of adverse effects
- (D) Recommend against routine use if no increased risk

Folic acid supplementation²⁴

- (A) 0.4 to 0.8 mg daily for women capable of conceiving

Statins for primary prevention of CVD²⁵

- (B) Recommend low- to moderate-dose statin therapy in patients meeting all three criteria:
 - (1) 40 to 75 years of age
 - (2) Dyslipidemia, diabetes, hypertension, or smoker
 - (3) 10-year CVD risk of 10% or greater
- (C) Consider low- to moderate-dose statin therapy in appropriate candidates meeting the first two criteria but with a 10-year CVD risk of 7.5% to 10%
- (I) IETRFOA initiating statin therapy after 75 years of age for primary prevention

Fall prevention in community-dwelling older adults²⁶

- (B) Recommend exercise interventions for individuals 65 years and older at increased risk of falls
- (C) Recommend multifactorial interventions for appropriate individuals 65 years and older; see Clinical Considerations in original recommendation statement for patient selection
- (D) Recommend against vitamin D supplementation for fall prevention

Counseling to prevent sexually transmitted infection²⁷

- (B) Recommend counseling to prevent sexually transmitted infection for adolescents and adults at increased risk

Counseling to promote healthy diet and physical activity²⁸

- (B) Recommend that patients with other CVD risk factor(s) who are overweight or obese be offered or referred for intensive behavioral counseling

Counseling for skin cancer prevention²⁹

- (B) Recommend counseling fair-skinned patients six months to 24 years of age about minimizing ultraviolet radiation
- (C) Recommend selectively counseling fair-skinned patients older than 24 years about minimizing exposure to ultraviolet radiation
- (I) IETRFOA counseling adults about skin self-examination

Counseling to promote healthy weight gain in pregnancy³⁰

- (B) Offer behavioral counseling interventions to promote health weight gain and to prevent excessive weight gain to all pregnant women

Grade C Recommendations:

- Initiation of low-dose aspirin for adults 40 to 59 years of age who have a 10% or greater 10-year CVD risk³¹
- Physical activity and healthy diet counseling to reduce cardiovascular risk in adults without obesity or known CVD risk factors³²
- Prostate cancer screening with prostate-specific antigen testing in men 55 to 69 years of age after shared decision-making³³

Grade D Recommendations:

- Bacteriuria (asymptomatic) screening in nonpregnant adults³⁴
- Beta carotene or vitamin E supplementation for CVD or cancer risk reduction³⁵
- Carotid artery stenosis screening³⁶
- CVD screening with resting or exercise electrocardiography in low-risk patients³⁷
- Chronic obstructive pulmonary disease screening with spirometry³⁸
- Hormone therapy for the primary prevention of chronic conditions in postmenopausal persons³⁹
- Genital herpes screening, serologic screening⁴⁰
- Initiation of low-dose aspirin for adults 60 years or older⁴¹
- Ovarian cancer screening⁴¹
- Pancreatic cancer screening⁴²
- Prostate cancer screening with prostate-specific antigen testing in men 70 years and older³³
- Testicular cancer screening⁴³
- Thyroid cancer screening⁴⁴
- Vitamin D (≤ 400 IU) and calcium ($\leq 1,000$ mg) supplementation daily for primary prevention of fracture in postmenopausal women⁴⁵

Grade I Statements:

- Atrial fibrillation screening⁴⁶
- Bladder cancer screening⁴⁷
- Celiac disease screening⁴⁸
- CVD screening in patients with nontraditional risk factors⁴⁹
- CVD screening with resting or exercise electrocardiography in intermediate- to high-risk patients³⁷
- Chronic kidney disease screening⁵⁰
- Cognitive impairment screening in older adults⁵¹
- Eating disorders screening⁵²
- Gynecologic condition screening with pelvic examination⁵³
- Hearing loss screening in older adults⁵⁴
- Impaired visual acuity screening in older adults⁵⁵
- Multivitamin, single nutrient, or paired nutrients for CVD or cancer risk reduction (beta carotene and vitamin E, as above)³⁵
- Obstructive sleep apnea screening⁵⁶
- Oral cancer screening⁵⁷
- Peripheral artery disease and CVD risk screening with ankle-brachial index⁵⁸

continues

CHD = coronary heart disease; CVD = cardiovascular disease; FRAX = Fracture Risk Assessment; IETRFOA = insufficient evidence to recommend for or against; PrEP = preexposure prophylaxis; USPSTF = U.S. Preventive Services Task Force.

Adult Preventive Health Care Schedule: Recommendations from the USPSTF *(continued)*

Primary open-angle glaucoma screening⁵⁹

Primary prevention of fractures with vitamin D and calcium supplementation (alone or combined; dose unspecified) in men or premenopausal women, and in postmenopausal women with daily dosages > 400 IU of vitamin D and > 1,000 mg of calcium⁴⁵

Skin cancer screening⁶⁰

Suicide risk screening⁶¹

Suicide risk screening for children and adolescents⁶⁵

Thyroid dysfunction screening⁶²

Vitamin D deficiency screening in community-dwelling nonpregnant adults⁶³

References

- Curry SJ, Krist AH, Owens DK, et al. Screening and behavioral counseling interventions to reduce unhealthy alcohol use in adolescents and adults: U.S. Preventive Services Task Force recommendation statement. *JAMA*. 2018;320(18):1899-1909.
- Krist KH, Davidson KW, Mangione CM, et al. Screening for unhealthy drug use: US Preventive Services Task Force recommendation statement. *JAMA*. 2020;323(22):2301-2309.
- Siu AL, Bibbins-Domingo K, Grossman DC, et al. Screening for depression in adults: U.S. Preventive Services Task Force recommendation statement. *JAMA*. 2016;315(4):380-387.
- Krist AH, Davidson KW, Mangione CM, et al. Screening for hypertension in adults: US Preventive Services Task Force reaffirmation recommendation statement. *JAMA*. 2021;325(16):1650-1656.
- Curry SJ, Krist AH, Owens DK, et al. Behavioral weight loss interventions to prevent obesity-related morbidity and mortality in adults: U.S. Preventive Services Task Force recommendation statement. *JAMA*. 2018;320(11):1163-1171.
- Krist AH, Davidson KW, Mangione CM, et al. Interventions for tobacco smoking cessation in adults, including pregnant persons: US Preventive Services Task Force recommendation statement. *JAMA*. 2021;325(3):265-279.
- Owens DK, Davidson KW, Krist AH, et al. Screening for hepatitis C virus infections in adolescents and adults: U.S. Preventive Services Task Force recommendation statement. *JAMA*. 2020;323(10):970-975.
- Owens DK, Davidson KW, Krist AH, et al. Screening for HIV infection: U.S. Preventive Services Task Force recommendation statement. *JAMA*. 2019;321(23):2326-2336.
- Krist AH, Davidson KW, Mangione CM, et al. Screening for hepatitis B virus infection in adolescents and adults: US Preventive Services Task Force recommendation statement. *JAMA*. 2020;324(23):2415-2422.
- Mangione CM, Barry MJ, Nicholson WK, et al. Screening for syphilis infection in nonpregnant adolescents and adults: US Preventive Services Task Force recommendation statement. *JAMA*. 2022;328(12):1243-1249.
- Mangione CM, Barry MJ, Nicholson WK, et al. Screening for latent tuberculosis infection in adults: US Preventive Services Task Force recommendation statement. *JAMA*. 2023;329(17):1487-1494.
- Owens DK, Davidson KW, Krist AH, et al. Risk assessment, genetic counseling, and genetic testing for *BRCA*-related cancer: U.S. Preventive Services Task Force recommendation statement [published correction appears in *JAMA*. 2918(322(18)):1830]. *JAMA*. 2019;322(7):652-665.
- Davidson KW, Barry MJ, Mangione CM, et al. Screening for chlamydia and gonorrhea: US Preventive Services Task Force Recommendation Statement. *JAMA*. 2021;326(10):949-956.
- Curry SJ, Krist AH, Owens DK, et al. Screening for intimate partner violence, elder abuse, and abuse of vulnerable adults: U.S. Preventive Services Task Force final recommendation statement. *JAMA*. 2018;320(16):1678-1687.
- Curry SJ, Krist AH, Owens DK, et al. Screening for cervical cancer: U.S. Preventive Services Task Force recommendation statement. *JAMA*. 2018;320(7):674-686.
- Davidson KW, Barry MJ, Magione CM, et al. Screening for prediabetes and type 2 diabetes: US Preventive Services Task Force recommendation statement. *JAMA*. 2021;326(8):736-743.
- Davidson KW, Barry MJ, Mangione CM, et al. Screening for colorectal cancer: US Preventive Services Task Force recommendation statement [published correction appears in *JAMA*. 2021;326(8):773]. *JAMA*. 2021;325(19):1965-1977.
- Siu AL. Screening for breast cancer: U.S. Preventive Services Task Force recommendation statement [published correction appears in *Ann Intern Med*. 2016;164(6):448]. *Ann Intern Med*. 2016;164(4):279-296.
- Krist AH, Davidson KW, Mangione CM, et al. Screening for lung cancer: US Preventive Services Task Force recommendation statement. *JAMA*. 2021;325(10):962-970.
- Curry SJ, Krist AH, Owens DK, et al. Screening for osteoporosis to prevent fractures: U.S. Preventive Services Task Force recommendation statement. *JAMA*. 2018;319(24):2521-2531.
- Owens DK, Davidson KW, Krist AH, et al. Screening for abdominal aortic aneurysm: US Preventive Services Task Force recommendation statement. *JAMA*. 2019;322(22):2211-2218.
- Owens DK, Davidson KW, Krist AH, et al. Preexposure prophylaxis for the prevention of HIV infection: U.S. Preventive Services Task Force recommendation statement. *JAMA*. 2019;321(22):2203-2213.
- Owens DK, Davidson KW, Krist AH, et al. Medication use to reduce risk of breast cancer: US Preventive Services Task Force recommendation statement. *JAMA*. 2019;322(9):857-867.
- Bibbins-Domingo K, Grossman DC, Curry SJ, et al. Folic acid supplementation for the prevention of neural tube defects: US Preventive Services Task Force recommendation statement. *JAMA*. 2017;317(2):183-189.
- Mangione CM, Barry MJ, Nicholson WK, et al. Statin use for the primary prevention of cardiovascular disease in adults: US Preventive Services recommendation statement. *JAMA*. 2022;328(8):746-753.
- Grossman DC, Curry SJ, Owens DK, et al. Interventions to prevent falls in community-dwelling older adults: US Preventive Services Task Force recommendation statement. *JAMA*. 2018;319(16):1696-1704.
- Krist AH, Davidson KW, Mangione CM, et al. Behavioral counseling interventions to prevent sexually transmitted infections: US Preventive Services Task Force recommendation statement. *JAMA*. 2020;324(7):674-681.
- Krist AH, Davidson KW, Mangione CM, et al. Behavioral counseling interventions to promote a healthy diet and physical activity for cardiovascular disease prevention in adults with cardiovascular risk factors: US Preventive Services Task Force recommendation statement. *JAMA*. 2020;324(20):2069-2075.
- Grossman DC, Curry SJ, Owens DK, et al. Behavioral counseling to prevent skin cancer: US Preventive Services Task Force recommendation statement. *JAMA*. 2018;319(11):1134-1142.
- Davidson KW, Barry MJ, Mangione CM, et al. Behavioral counseling interventions for healthy weight and weight gain in pregnancy: US Preventive Services Task Force recommendation statement. *JAMA*. 2021;325(20):2087-2093.
- Davidson KW, Barry MJ, Mangione CM, et al. Aspirin use to prevent cardiovascular disease: US Preventive Services Task Force recommendation statement. *JAMA*. 2022;327(16):1577-1584.
- Mangione CM, Barry MJ, Nicholson WK, et al. Behavioral counseling interventions to promote a healthy diet and physical activity for cardiovascular disease prevention in adults without cardiovascular risk factors: US Preventive Services Task Force recommendation statement. *JAMA*. 2022;328(4):367-374.
- Grossman DC, Curry SJ, Owens DK, et al. Screening for prostate cancer: US Preventive Services Task Force recommendation statement [published correction appears in *JAMA*. 2018;319(23):2443]. *JAMA*. 2018;319(18):1901-1913.
- Owens DK, Davidson KW, Krist AH, et al. Screening for asymptomatic bacteriuria in adults: US Preventive Services Task Force recommendation statement. *JAMA*. 2019;322(12):1188-1194.
- Mangione CM, Barry MJ, Nicholson WK, et al. Vitamin, mineral, and multivitamin supplementation to prevent cardiovascular disease and cancer: US Preventive Services Task Force recommendation statement. *JAMA*. 2022;327(23):2326-2333.
- Krist AH, Davidson KW, Mangione CM, et al. Screening for asymptomatic carotid artery stenosis: US Preventive Services Task Force recommendation statement. *JAMA*. 2021;325(5):476-481.
- Curry SJ, Krist AH, Owens DK, et al. Screening for cardiovascular disease risk with electrocardiography: US Preventive Services Task Force recommendation statement. *JAMA*. 2018;319(22):2308-2314.

Adult Preventive Health Care Schedule: Recommendations from the USPSTF *(continued)*

38. Mangione CM, Barry MJ, Nicholson WK, et al. Screening for chronic obstructive pulmonary disease: US Preventive Services Task Force reaffirmation recommendation statement. *JAMA*. 2022;327(18):1806-1811.
39. Mangione CM, Barry MJ, Nicholson WK, et al. Hormone therapy for the primary prevention of chronic conditions in postmenopausal persons: US Preventive Services Task Force recommendation statement. *JAMA*. 2022;328(17):1740-1746.
40. Mangione CM, Barry MJ, Nicholson WK, et al. Serologic screening for genital herpes infection: US Preventive Services Task Force recommendation statement. *JAMA*. 2023;329(6):502-507.
41. Grossman DC, Curry SJ, Owens DK, et al. Screening for ovarian cancer: US Preventive Services Task Force recommendation statement. *JAMA*. 2018;319(6):588-594.
42. Owens DK, Davidson KW, Krist AH, et al. Screening for pancreatic cancer. US Preventive Services Task Force reaffirmation recommendation statement. *JAMA*. 2019;322(5):438-444.
43. Screening for testicular cancer: U.S. Preventive Services Task Force reaffirmation recommendation statement. *Ann Intern Med*. 2011;154(7):483-486.
44. Bibbins-Domingo K, Grossman DC, Curry SJ, et al. Screening for thyroid cancer: US Preventive Services Task Force recommendation statement. *JAMA*. 2017;317(18):1882-1887.
45. Grossman DC, Curry SJ, Owens DK, et al. Vitamin D, calcium, or combined supplementation for the primary prevention of fractures in community-dwelling adults: US Preventive Services Task Force recommendation statement. *JAMA*. 2018;319(15):1592-1599.
46. Davidson KW, Barry MJ, Mangione CM, et al. Screening for atrial fibrillation: U.S. Preventive Services Task Force recommendation statement. *JAMA*. 2022;327(4):360-367.
47. Moyer VA. Screening for bladder cancer: U.S. Preventive Services Task Force recommendation statement [published correction appears in *Ann Intern Med*. 2011;155(6):408]. *Ann Intern Med*. 2011;155(4):246-251.
48. Bibbins-Domingo K, Grossman DC, Curry SJ, et al. Screening for celiac disease: US Preventive Services Task Force recommendation statement. *JAMA*. 2017;317(12):1252-1257.
49. Curry SJ, Krist AH, Owens DK, et al. Risk assessment for cardiovascular disease with nontraditional risk factors: US Preventive Services Task Force recommendation statement. *JAMA*. 2018;320(3):272-280.
50. Moyer VA. Screening for chronic kidney disease: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med*. 2012;157(8):567-570.
51. Owens DK, Davidson KW, Krist AH, et al. Screening for cognitive impairment in older adults: US Preventive Services Task Force recommendation statement. *JAMA*. 2020;323(8):757-763.
52. Davidson KW, Barry MJ, Mangione CM, et al. Screening for eating disorders in adolescents and adults: US Preventive Services Task Force recommendation statement. *JAMA*. 2022;327(11):1061-1067.
53. Bibbins-Domingo K, Grossman DC, Curry SJ, et al. Screening for gynecologic conditions with pelvic examination: US Preventive Services Task Force recommendation statement. *JAMA*. 2017;317(9):947-953.
54. Krist AH, Davidson KW, Mangione CM, et al. Screening for hearing loss in older adults: US Preventive Services Task Force recommendation statement. *JAMA*. 2021;325(12):1196-1201.
55. Mangione CM, Barry MJ, Nicholson WK, et al. Screening for impaired visual acuity in older adults: US Preventive Services Task Force recommendation statement. *JAMA*. 2022;327(21):2123-2128.
56. Mangione CM, Barry MJ, Nicholson WK, et al. Screening for obstructive sleep apnea in adults: US Preventive Services Task Force recommendation statement. *JAMA*. 2022;328(19):1945-1950.
57. Moyer VA. Screening for oral cancer: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med*. 2013;160(1):55-60.
58. Curry SJ, Krist AH, Owens DK, et al. Screening for peripheral artery disease and cardiovascular disease risk assessment with the ankle-brachial index: US Preventive Services Task Force recommendation statement. *JAMA*. 2018;320(2):177-183.
59. Mangione CM, Barry MJ, Nicholson WK, et al. Screening for primary open-angle glaucoma: US Preventive Services Task Force recommendation statement. *JAMA*. 2022;327(20):1992-1997.
60. Mangione CM, Barry MJ, Nicholson WK, et al. Screening for skin cancer: US Preventive Services Task Force recommendation statement. *JAMA*. 2023;329(15):1290-1295.
61. LeFevre ML. Screening for suicide risk in adolescents, adults, and older adults in primary care: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med*. 2014;160(10):719-726.
62. LeFevre ML. Screening for thyroid dysfunction: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med*. 2015;162(9):641-650.
63. Krist AH, Davidson KW, Mangione CM, et al. Screening for vitamin D deficiency in adults: US Preventive Services Task Force recommendation statement. *JAMA*. 2021;325(14):1436-1442.